

12 Month - Ryegrass Forage Value List



Evaluation date: Jan 2020

Lower Nth. Island



Cultivars are sorted by star rating and then alphabetically. Note:

- The short term ryegrasses are sown by dairy farmers for 12 month production.
- The FVI for 12 month ryegrasses is a combination of seasonal dry matter production performance and economic values only.
- WE is without endophyte or also known as nil endophyte.
- 12 month options include Hybrid and Italian ryegrasses.

Filtered by: LOWER NORTH ISLAND, 12 MONTH/AR37/NEA/NEA2/SE/WE, TETRAPLOID, MID/LATE/VERY LATE

FVI ¹ (Star rating)	FVI Star Rating (\$/ha)	Cultivar	Performance Values (1-5 rating) Seasonal dry matter (DM)					Other Cultivar Information				
			EST	Winter	Early Spring	Late Spring	Summer	Endo ⁶	Ploidy ⁷	HD ⁸	Marketer	Conf ⁹
★★★★★	\$668 to \$889	■ SHOGUN NEA	2	3	5	5	5	NEA	Tetraploid	VL	Barenbrug	10+
★★★★	\$448 to \$668	● FEAST II WE	3	3	2	2	4	WE	Tetraploid	L	PGG Wrightson Seeds	10+
★★★★	\$448 to \$668	● LUSH AR37	5	4	1	1	4	AR37	Tetraploid	L	PGG Wrightson Seeds	10+
★★★	\$227 to \$448	● MONA WE	3	3	3	3	3	WE	Tetraploid	VL	DLF Seeds	7.4
★	-\$214 to \$7	● MOATA WE	1	1	1	1	1	WE	Tetraploid	L	Common	10+

¹5 = Top rank, 1 = bottom rank, ²EST=establishment dry matter production (Mar-May), Winter = Winter dry matter production (June-July), Early Spring = Early spring dry matter production (Aug-Sept), Late Spring = Late spring dry matter production (Oct-Nov), Summer = Summer dry matter production (Dec-Feb), ³Endophyte, WE is without endophyte, ⁴Ploidy (D=Diploid, T=Tetraploid), ⁵Heading date (MS=mid season, L=Late, VL=Very late), ⁶Confidence (based on number of DM trials in the regions weighted by the DM trait correlations).

For more information visit www.dairynz.co.nz/fvi.

DairyNZ Limited and its agents and employees ("DairyNZ") provide no assurance or warranty as to the accuracy, completeness or reliability of information in the Forage Value Index or at dairynz.co.nz/fvi. DairyNZ has no liability for any reliance on that information.